

# UK Maths Trust

## MATHEMATICAL COMPETITION FOR GIRLS

© 2025 UK Mathematics Trust

supported by  Jane Street® 

## SOLUTIONS

Although in this paper you were only required to provide numerical answers, we present full solutions, including all the reasoning leading to those answers. These will be helpful if you ever enter an olympiad which requires full solutions.

The answers that needed to be entered on the answer sheets are given at the start of each solution.

These are polished solutions and do not illustrate the process of failed ideas and rough work by which candidates may arrive at their own solutions. They are not intended to be the ‘best’ possible solutions; in some cases we have suggested alternatives, but readers may come up with other equally good ideas.

Enquiries about the Mathematical Competition for Girls should be sent to:

*challenges@ukmt.org.uk*

[www.ukmt.org.uk](http://www.ukmt.org.uk)

**1.** Before you start this question, expand and simplify  $(a + b)^3$ .

For each part of the question, enter your answer as follows. The first digit should be the number of digits of your answer, and the second two digits should be the sum of the digits of your answer. For example, if your answer is 2025, you should enter 409 on the answer sheet (2025 has 4 digits, and the sum of its digits is 09).

- (a) Find the cube root of 1030301. [3 marks]  
 (b) Find the square root of 16008001. [3 marks]  
 (c) Find the fourth root of 16096216216081. [4 marks]

Answer: (a) 302 (b) 405 (c) 405

**COMMENTARY**

This question is about using algebraic identities to evaluate numerical expressions. For example, you may be familiar with using the difference of two squares to quickly evaluate expressions such as  $998^2 - 4$ : you write it as  $998^2 - 2^2 = (998 - 2)(998 + 2)$  from which you can easily see that the answer is 996000.

For the first part, the key is to look at the expanded form of  $(a + b)^3$  and write the number 1030301 as a sum in which each part corresponds to one term in the expansion.


You will need different expansions for the other two parts.

- (a) Using the expansion of  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$  with  $a = 100$  and  $b = 1$  we can see that  $101^3 = 1000000 + 30000 + 300 + 1 = 1030301$ , so the required cube root is 101.  
 (b) Using the expansion of  $(a + b)^2 = a^2 + 2ab + b^2$  with  $a = 4000$ ,  $b = 1$  we get that  $4001^2 = 16000000 + 8000 + 1 = 16008001$ , so the required square root is 4001.  
 (c) Using  $(a + b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4$ , we find that

$$\begin{aligned} 2003^4 &= 16 \times 10^{12} + 4(8 \times 10^9 \times 3) + 6(4 \times 10^6 \times 9) + 4(2 \times 10^3 \times 27) + 81 \\ &= 16 \times 10^{12} + 96 \times 10^9 + 216 \times 10^6 + 216 \times 10^3 + 81 \end{aligned}$$

so the required fourth root is 2003.

2. Tom has a large supply of two types of dominoes, Type A and Type B. Type B dominoes can be rotated  $180^\circ$  so that the grey square is on top.



Tom wants to select three dominoes and place them next to each other to create a  $2 \times 3$  rectangle (so the dominoes remain vertical, as shown above). He wants both top and bottom rows of his rectangle to contain at least one square of each colour.

- (a) (i) How many sequences can Tom make which contain exactly one Type A domino? [2 marks]

- (ii) How many sequences of three dominoes can he make in total? [2 marks]

- (b) Each of the white squares on Type B dominoes has a whole number between 1 and 6 (inclusive) written on it. Grey squares have no numbers on them. There are several copies of each numbered Type B domino.

(The numbers do not change when a Type B domino is rotated; for example a 6 does *not* become a 9.)

Tom wants to select and arrange three dominoes as before (with both top and bottom rows containing at least one square of each colour, and using dominoes of either type), but now he also wants the numbers on the top row to add up to 6 and the numbers on the bottom row to add up to 6.

In how many ways can he do this? [6 marks]

Answer: (a)(i) 006 (ii) 012 (b) 036

#### COMMENTARY

The key to counting questions is to organise your work systematically, making sure you consider all possible cases and do not include any case more than once.

If there are not too many options it is fine to list them all, as long as you explain how you have ensured that you have not missed any. It is also helpful to realise when you can group the options in some way, and then only list some of the groups. For example, in part (a)(i), there are three positions where the Type A domino can go. So you can put it in the first position, list all possible options for the other two dominoes, and then multiply that number by three. (Note that, in the solution below, we have taken a different approach.)

The structure of the question strongly suggests that for sub-part (a)(ii) you should consider two separate cases: one including a Type A domino and one without it. Note that you need to explain why those are the only cases (i.e. why you can't have more than one Type A domino).

Part (b) can be linked to part (a). For each arrangement from part (a), think about how you can number the white squares in order to satisfy the conditions.

- (a) (i) When using a Type A domino, Tom must also use two Type B dominoes, one in each orientation, in order to have one white square in each row. He is therefore using three different-looking dominoes, and they can be arranged in  $3 \times 2 \times 1 = 6$  ways.
- (ii) Call the two orientations of the Type B domino (as shown in the diagram)  $B_1$  and  $B_2$ . Tom must use at least one of each of these two types, otherwise one of the rows would contain only grey squares.

This means that Tom can use the following sets of dominoes: one of each type; one  $B_1$  and two  $B_2$ s; or one  $B_2$  and two  $B_1$ s.

The first case gives the six sequences described in part (i). The second and the third case give the same number of sequences, so we only need to look at the second case.

The single  $B_1$  domino can go in any of the three places, with the remaining places being taken up by the two  $B_2$  dominoes. Hence there are three sequences for the second case, and another three for the third case.

This gives the total of  $6 + 3 + 3 = 12$  possible sequences.

- (b) We consider the three cases described in the previous part. For each possible sequence of the dominoes, we count how many ways there are to number them.

In the case with one of each type of domino, each row contains only one white square, so they must both have the number 6 written on them. There is only one possible numbering for each of the six sequences, giving six possible ways for this case.

In the case with one  $B_1$  and two  $B_2$  dominoes, the  $B_1$  domino must have the number 6 on it, but the two  $B_2$  dominoes can have one of the following five pairs:  $(1, 5)$ ,  $(2, 4)$ ,  $(3, 3)$ ,  $(4, 2)$  or  $(5, 1)$ . So for each of the three sequences of dominoes, there are five ways to select the numbers. This gives  $3 \times 5 = 15$  possibilities for the second case.

The third case is the same as the second case. So the total number of possibilities is  $6 + 15 + 15 = 36$ .

3. Consider the number  $M = 99\dots 99$  which consists of several digit nines. A single division sign is placed between two adjacent digits of  $M$  and the resulting calculation is evaluated to produce a whole number  $N$ .

(a) In the case when  $M$  has nine digits,

(i) How many possible values can  $N$  take? [1 mark]

(ii) How many digits does the smallest possible value of  $N$  have? [2 marks]

(b) In the case when  $M$  has 2025 digits,

(i) How many possible values can  $N$  take? [3 marks]

(ii) How many digits does the smallest possible value of  $N$  have? [4 marks]

Answer: (a)(i) 002 (ii) 004 (b)(i) 014 (ii) 676

#### COMMENTARY

In a question like this it is sensible to start by looking at some small cases, which is what part (a) is encouraging you to do. You can look at examples with even fewer nines to develop and test your ideas.

The key is to come up with a simple way to do the division.

For example, if we want to divide a number with six nines by a number with two nines, we can write the first number as  $990000 + 9900 + 99$ , so dividing it by 99 gives  $10000 + 100 + 1 = 10101$ . On the other hand, if the first number has five nines, it equals  $99000 + 990 + 9$ , which is not a multiple of 99.

Trying some examples like this will hopefully lead you to an idea you can apply in both parts of the question. You should note both when the division gives a whole number, and what the form of the answer is.

Write the division as  $A \div B$ , where  $A$  consists of  $a$  nines and  $B$  consists of  $b$  nines. We are going to show that the calculation produces a whole number if, and only if,  $a$  is a multiple of  $b$ . We are then going to apply that idea to both parts of the question.

We must have  $A \geq B$  for the division to produce a whole number so  $a \geq b$ . Write  $a = kb + r$ , where  $k \geq 1$  and  $0 \leq r < b$ . This means that we can split  $A$  into  $k$  blocks of  $b$  nines, going from left to right, with  $r$  nines left over.

Each block of  $b$  nines represents a number which is  $B$  followed by some zeros. It is therefore divisible by  $B$ . The remaining  $r$  nines represent a number which is smaller than  $B$ , so dividing  $A$  by  $B$  leaves a remainder which is a number with  $r$  nines.

Therefore, the only way that  $A \div B$  can be a whole number is if  $r = 0$ , in which case  $a$  is a multiple of  $b$ .

(a) The possible ways to write 9 as  $a + b$  with  $a \geq b \geq 1$  are  $8 + 1$ ,  $7 + 2$ ,  $6 + 3$  and  $5 + 4$ . Out of those,  $a$  is a multiple of  $b$  only in the cases  $a = 8$ ,  $b = 1$  and  $a = 6$ ,  $b = 3$ . Therefore  $N$

can take two possible values.

The smallest possible value of  $N$  is when  $a$  and  $b$  are closest together since this minimises  $A$  and maximises  $B$ . In this case, it is  $999999 \div 999 = 1001$ , which has 4 digits.

- (b) We need to write  $2025 = a + b$  where  $a \geq b \geq 1$  and  $a$  is a multiple of  $b$ . Each of these pairs of  $a$  and  $b$  will give a different value of the whole number  $N$ , with the smallest  $N$  corresponding to the pair in which  $a$  and  $b$  are closest together, meaning that  $b$  is as large as possible.

Writing  $a = kb$ , with  $k \geq 1$ , we have  $2025 = kb + b = (k + 1)b$ . This means that  $(k + 1)$  and  $b$  are factors of 2025 with  $(k + 1) \geq 2$ .

We know that 2025 has 15 factors. However, one of them is 1, which is not a possible value of  $(k + 1)$ . Hence, there are 14 possible values of  $N$ .

The smallest value of  $N$  occurs when  $b$  is as large as possible. This is the case when  $(k + 1) = 3$ ,  $b = 675$ , resulting in  $a = kb = 1350$ . In the division  $A \div B$ ,  $A$  consists of two blocks of 675 nines, so  $N$  has 676 digits (a 1 followed by 674 zeros followed by another 1).

#### NOTE

The proof that  $b$  needs to divide  $a$  can also be written using algebra, instead of thinking about blocks of nines.

Note that the number with  $a$  nines can be written as  $10^a - 1$ . It is well known that  $x^k - 1$  has a factor of  $x - 1$  for all positive integers  $k$ .

If  $a = kb + r$ , then

$$\begin{aligned} 10^a - 1 &= 10^{kb} \times 10^r - 1 \\ &= 10^r (10^{kb} - 1) + (10^r - 1) \\ &= 10^r ((10^b)^k - 1) + (10^r - 1) \end{aligned}$$

The first term has a factor of  $10^b - 1$ , which is our number  $B$ . Since  $r < b$ , the second term is smaller than  $B$ , so that is the remainder of the division and equals zero if, and only if,  $r = 0$ .

The full result mentioned above says that  $x^k - 1 = (x - 1)(x^{k-1} + x^{k-2} + \dots + x + 1)$ , and this tells you the form of the resulting integer  $N$ .

4. (a) Positive (non-zero) whole numbers  $a$  and  $b$  satisfy  $(a + b)(a - b) = 45$ .
- (i) How many possible values of  $a$  are there? [2 marks]
- (ii) What is the smallest possible value of  $a$ ? [2 marks]
- (b) Priya and Rhia each create a sequence of positive integers.
- Priya starts with 1000 and adds consecutive odd numbers, so that her sequence begins 1000, 1001, 1004, 1009.
- Rhia starts with 3025 and also adds consecutive odd numbers.
- (i) The number 3025 appears in both sequences. The next smallest number that appears in both sequences is a four-digit number ' $3abc$ '. Enter the digits ' $abc$ ' on your answer sheet. [3 marks]
- (ii) How many numbers (including 3025) appear in both sequences? [3 marks]

Answer: (a)(i) 003 (ii) 007 (b)(i) 601 (ii) 008

#### COMMENTARY

This is another question where part (a) gives you a strong hint for how to approach part (b): you are looking to write an equation which you can factorise, and then look for factors of some number.

You may already know that consecutive odd numbers add up to a square number (for example,  $1 + 3 + 5 + 7 = 16$ ). Even if you don't, you should be able to spot this pattern by continuing Priya's sequence for a few more terms. This is an example of a known result that could be stated without proof, even in a full solution question.

Note that in the solution below we use two different letters,  $n$  and  $k$ , for the general terms of the two sequences. A common mistake is to use the same letter, but that would mean that the equal terms are always in the same position, which is clearly not the case.

- (a) Both  $(a + b)$  and  $(a - b)$  must be factors of 45. Furthermore, as  $b$  is positive,  $(a + b)$  must be the larger factor. We therefore have the following possibilities:

$a + b$	$a - b$	$a$
45	1	23
15	3	9
9	5	7

The possible values of  $a$  are 7, 9 and 23.

- (b) Adding consecutive odd numbers, starting from 1, gives consecutive square numbers. Therefore the  $n$ th term of Priya's sequences is  $1000 + n^2$  and the  $k$ th term of Rhia's sequence is  $3025 + k^2$ , where  $n, k \geq 0$  (the "0th term" corresponds to the starting number).

For a number to appear in both sequences, we need

$$\begin{aligned}1000 + n^2 &= 3025 + k^2 \\ \iff n^2 - k^2 &= 2025 \\ \iff (n - k)(n + k) &= 2025\end{aligned}$$

This shows that both  $(n - k)$  and  $(n + k)$  need to be factors of 2025. Since  $k \geq 0$ , we must have  $n + k \geq n - k$ . Note that we need not consider negative factors because  $n$  and  $k$  are both positive, so it is not possible for both  $n + k$  and  $n - k$  to be negative.

The solutions therefore correspond to the factor pairs of 2025, with  $n + k$  being the larger one. Note that 2025 is odd, giving  $n - k$ ,  $n + k$  both odd, so  $n$  and  $k$  are indeed both integers for each factor pair.

- (i) Smaller numbers in the sequence correspond to the smaller values of  $k$ . The value of  $k$  will be the smallest when the two factors of 2025 are closest together.


Note that  $n - k = n + k = 45$  is allowed, as this corresponds to  $k = 0, n = 45$ , giving  $1000 + 45^2 = 3025 + 0^2 = 3025$  as the smallest number which appears in both sequences.

The next factor pair where the factors are closest to each other is  $27 \times 75$ . This gives  $n - k = 27$  and  $n + k = 75$  so  $k = 24$ . The second smallest number that appears in both sequences is  $3025 + 24^2 = 3601$ .

- (ii) Since 2025 has fifteen factors, it has eight factor pairs, including  $45 \times 45$ . Hence there are eight numbers which appear in both sequences.



5. The diagram shows triangle  $ABC$  with side lengths  $AB = 85$ ,  $BC = 160$  and  $CA = 103$  units. The bisector of angle  $ABC$  intersects the side  $AC$  at point  $D$ . The line through  $A$  perpendicular to  $BD$  intersects  $BD$  at  $M$  and  $BC$  at  $P$ .



- (a) Which of the following statements are correct?

- (i)  $D$  is the midpoint of  $AC$ .
- (ii)  $M$  is the midpoint of  $AP$ .
- (iii)  $P$  is the midpoint of  $BC$ .

Enter 1 for each correct statement and 0 for each incorrect statement. For example, if you think that statements (i) and (ii) are correct, you should enter 110. [2 marks]

- (b) State the length of  $PC$ . [2 marks]

- (c) The line through  $A$  perpendicular to the bisector of angle  $ACB$  intersects it at  $N$ .

Find the length of  $MN$ . [6 marks]

Answer: (a) 010 (b) 075 (c) 014

#### COMMENTARY

This is a geometry problem where each part builds on the previous one.

In part (a), the goal is to show lengths  $AM$  and  $MP$  are equal. Since triangles  $AMB$  and  $PMB$  share a side and have a right angle at  $M$ , it is natural to try to prove the two triangles are congruent. There are several congruence rules (ASA, SAS, SSS, RHS), so the task is to identify the one that is most useful here (in fact, multiple work).

For part (b), it helps to mark the given lengths on the diagram. We already know length  $BC$ , so finding length  $BP$  would allow us to work out length  $PC$ . This connects back to some of the work we did in part (a).

In the final part, the problem introduces a similar construction on the other side of the triangle. The results from parts (a) and (b) already give useful information about the diagram. The final step is to find a way to link the new segment  $MN$  to lengths we can compute using what we have established earlier.


It may be helpful to redraw the diagram to show only the information relevant to part (c), as we did in the solution below.

- (a) Statement (i) is false. The angle bisector does not in general meet the opposite side at its midpoint. (This only happens when the triangle is isosceles.)

Statement (ii) is true. Since the line  $BM$  bisects angle  $PBA$ , angles  $PBM$  and  $ABM$  are equal. Since  $AM$  is perpendicular to  $BM$ , angles  $BMP$  and  $BMA$  both equal  $90^\circ$ . Therefore, triangles  $BPM$  and  $BAM$  have two equal angles and share the side  $BM$ , so they are congruent (ASA), which means that  $MP = MA$ .

Statement (iii) is false, as will be shown in part (b) by calculating the length of  $PC$ .

- (b) From the congruence of triangles  $BPM$  and  $BAM$  it also follows that  $BP = BA = 85$ . Hence  $PC = BC - BP = 75$ .
- (c) Let the line  $AN$  intersect  $BC$  at  $Q$ . By the same argument as above,  $N$  is the midpoint of  $AQ$  and  $CQ = CA = 103$ . We can then draw the following diagram:



In triangle  $AQP$ ,  $N$  and  $M$  are the midpoints of the sides  $AQ$  and  $AP$ . Therefore the length of  $NM$  is half the length of  $QP$ . But  $QP = QC - PC = 103 - 75 = 28$ , so the required length of  $MN$  is 14.